



Figure caption: Schematic diagram illustrating the manner in which early tendinopathy may arise due to hypoxic injury. See p 309.

Editor

Tore K Kvien

Associate Editors

Francis Berenbaum
Hans Bijlsma
David Pissetsky
Dimitrios Boumpas
Gerd Burmester
Mary Crow
Iain McInnes

Editorial office

Annals of the Rheumatic Diseases
BMJ Publishing Group Ltd
BMA House
Tavistock Square
London WC1H 9JR, UK
T: +44 (0)20 7383 6250
F: +44 (0)20 7383 6668
E: ard@bmj.com
ISSN: 0003-4967 (print)
ISSN: 1468-2060 (online)
Impact Factor: 9.082

Disclaimer: ARD is owned and published by BMJ Publishing Group Ltd (a wholly owned subsidiary of the British Medical Association) and the European League Against Rheumatism. The owners grant editorial freedom to the Editor of ARD. ARD follows guidelines on editorial independence produced by the World Association of Medical Editors and the code on good publication practice of the Committee on Publication Ethics.

ARD is intended for medical professionals and is provided without warranty, express or implied. Statements in the journal are the responsibility of their authors and advertisers and not authors' institutions the BMJ Publishing Group, the European League Against Rheumatism or the BMA unless otherwise specified or determined by law. Acceptance of advertising does not imply endorsement.

To the fullest extent permitted by law, the BMJ Publishing Group shall not be liable for any loss, injury or damage resulting from the use of ARD or any information in it whether based on contract, tort, or otherwise. Readers are advised to verify any information they choose to rely on.

Copyright: © 2012 BMJ Publishing Group and European League Against Rheumatism. All rights reserved; no part of this publication may be reproduced stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying recording, or otherwise without prior permission

ARD is published by BMJ Publishing Group Ltd typeset by Newgen and printed in the UK on acid-free paper from sustainable forests by Buxton Press, Buxton

Annals of the Rheumatic Diseases (ISSN No: 0003-4967) is published monthly by BMJ Publishing Group and distributed in the USA by Mercury International Ltd. Periodicals postage paid at Rahway, NJ, USA. POSTMASTER: send address changes to Annals of the Rheumatic Diseases Mercury International Ltd, 365 Blair Road, Avenel NJ 07001, USA.

Editorial

- 159** Canes for knee osteoarthritis: is a randomised trial necessary?
K B Hagen

Review

- 161** The effect of biological agents on work participation in rheumatoid arthritis patients: a systematic review
M M ter Wee, W F Lems, H Usan, A Gulpen, A Boonen

Clinical and epidemiological research

- 172** Impact of cane use on pain, function, general health and energy expenditure during gait in patients with knee osteoarthritis: a randomised controlled trial
A Jones, P G Silva, A C Silva, M Colucci, A Tuffanin, J R Jardim, J Natour

- 180** Safety, tolerability, pharmacokinetics, pharmacodynamics and efficacy of the monoclonal antibody ASK8007 blocking osteopontin in patients with rheumatoid arthritis: a randomised, placebo controlled, proof-of-concept study
M J H Boumans, J G A Houbiers, P Verschueren, H Ishikura, R Westhovens, E Brouwer, B Rojkevich, S Kelly, M den Adel, J Isaacs, H Jacobs, J Gomez-Reino, G M Holtkamp, A Hastings, D M Gerlag, P P Tak

- 186** In early rheumatoid arthritis, patients with a good initial response to methotrexate have excellent 2-year clinical outcomes, but radiological progression is not fully prevented: data from the methotrexate responders population in the SWEFOT trial
H Rezaei, S Saevarsdottir, K Forslind, K Albertsson, H Wallin, J Bratt, S Ernestam, P Geborek, I F Petersson, R F van Vollenhoven

- 192** The prospective association between psychological distress and disease activity in rheumatoid arthritis: a multilevel regression analysis
C L Overman, E R Bossema, H van Middendorp, L Wijngaards-de Meij, S MM Verstappen, M Bulder, J WG Jacobs, J WJ Bijlsma, R Geenen

- 198** Efficacy of tocilizumab in patients with moderate to severe active rheumatoid arthritis and a previous inadequate response to disease-modifying antirheumatic drugs: the ROSE study
Y Yazici, J R Curtis, A Ince, H Baraf, R L Malamet, L L Teng, A Kavanaugh

- 206** Predicting low disease activity and remission using early treatment response to antitumour necrosis factor therapy in patients with rheumatoid arthritis: exploratory analyses from the TEMPO trial
J R Curtis, S Yang, L Chen, G S Park, B Bitman, B Wang, I Navarro-Millan, A Kavanaugh

- 213** Disability in rheumatoid arthritis in the era of biological treatments
E Krishnan, B Lingala, B Bruce, J F Fries

- 219** The association between smoking and the development of psoriatic arthritis among psoriasis patients
L Eder, S Shanmugarajah, A Thavaneswaran, V Chandran, C F Rosen, R J Cook, D D Gladman

- 225** A mixed treatment comparison of the efficacy of anti-TNF agents in rheumatoid arthritis for methotrexate non-responders demonstrates differences between treatments: a Bayesian approach
S Schmitz, R Adams, C D Walsh, M Barry, O FitzGerald

- 231** Biphasic emergence of active tuberculosis in rheumatoid arthritis patients receiving TNF α inhibitors: the utility of IFN γ assay
D-Y Chen, G-H Shen, Y-M Chen, H-H Chen, C-W Hsieh, J-L Lan

- 238** Undifferentiated arthritis characteristics and outcomes when applying the 2010 and 1987 criteria for rheumatoid arthritis
A Krabben, T W J Huizinga, A H M van der Helm-van Mil

- 242** Clinical manifestations of antiphospholipid syndrome (APS) with and without antiphospholipid antibodies (the so-called 'seronegative APS')
J L Rodriguez-Garcia, M L Bertolaccini, M J Cuadrado, G Sanna, O Ateka-Barrutia, M A Khamashia

- 245** The association of treatment response and joint damage with ACPA-status in recent-onset RA: a subanalysis of the 8-year follow-up of the BeSt study
M van den Broek, L Dirven, NB Klarenbeek, THE Molenaar, KH Han, PJS M Kerstens, TWJ Huizinga, BAC Dijkemans, CF Allaart

- 249** The pulmonary arterial hypertension quality enhancement research initiative: comparison of patients with idiopathic PAH to patients with systemic sclerosis-associated PAH
P J Clements, M Tan, V V McLaughlin, R J Oudiz, V F Tapson, R N Channick, L J Rubin, A Langer, on behalf of the Pulmonary Arterial Hypertension Quality Enhancement Research Initiative (PAH-QuERI) Investigators

Basic and translational research

- 253** The anti-inflammatory effects of sympathectomy in murine antigen-induced arthritis are associated with a reduction of Th1 and Th17 responses
M Ebbinghaus, M Gajda, M K Boettger, H-G Schaible, R Bräuer

MORE CONTENTS ►



This article has been chosen by the editor to be of special interest or importance and is freely available online.



Articles carrying the Unlocked Logo are freely available online under the BMJ Journals unlocked scheme. See <http://ard.bmj.com/info/unlocked.dtl>

C O P E | COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk



When you have finished with this please recycle it

- 262** Comparison of rheumatoid articular adipose and synovial tissue reactivity to proinflammatory stimuli: contribution to adipocytokine network
E Kontny, M Plebanczyk, B Lisowska, M Olszewska, P Maldyk, W Maslinski
- 268** The ACPA recognition profile and subgrouping of ACPA-positive RA patients
A Willemze, S Böhringer, R Knevel, EW N Levarht, G Stoecken-Rijsbergen, JJ Houwing-Duistermaat, A HM van der Helm-van Mil, T WJ Huizinga, REM Toes, LA Trouw
- 275** Transcriptome analysis reveals specific changes in osteoarthritis synovial fibroblasts
M J Del Rey, A Usategui, E Izquierdo, J D Cañete, F J Blanco, G Criado, J L Pablos
- 281** Articular inflammation is controlled by myeloid cell-derived interleukin 1 receptor antagonist during the acute phase of arthritis in mice
C Lamacchia, E Rodriguez, G Palmer, S Vigne, P Martin, D Talbot-Ayer, C A Seemayer, C Gabay
- 288** Infrapatellar fat pad of patients with end-stage osteoarthritis inhibits catabolic mediators in cartilage
Y M Bastiaansen-Jenniskens, S Clockaerts, C Feijt, A-M Zuurmond, V Stojanovic-Susulic, C Bräts, L de Clerck, J DeGroot, J AN Verhaar, M Kloppenburg, G JVM van Osch
- 295** Potential involvement of IL-22 and IL-22-producing cells in the inflamed salivary glands of patients with Sjögren's syndrome
F Ciccia, G Guggino, A Rizzo, A Ferrante, S Raimondo, A Giardina, F Dieli, G Campisi, R Alessandro, G Triolo
- 302** Hypoxia: a critical regulator of early human tendinopathy
N L Millar, J H Reilly, S C Kerr, A L Campbell, K J Little, W J Leach, B P Rooney, G A C Murrell, I B McInnes
-
- Letters**
- 311** Cyclophosphamide is a highly effective and safe induction therapy in chronic periaortitis: a long-term follow-up of 35 patients with chronic periaortitis
M Binder, M Uhl, T Wiech, F Kollert, J Thiel, J O Sass, U A Walker, H H Peter, K Warnatz
- 313** Investigation of *IL1*, *VEGF*, *PPARG* and *MEFV* genes in psoriatic arthritis susceptibility
J Bowes, P Ho, E Flynn, S Salah, N McHugh, O FitzGerald, J Packham, A W Morgan, P S Helliwell, I N Bruce, A Barton
- 314** Association study of *BAK1* gene polymorphisms in Spanish rheumatoid arthritis and systemic lupus erythematosus cohorts
L-M Diaz-Gallo, S Garcia, N Ortego-Centeno, J Jiménez-Alonso, J Sánchez-Román, E de Ramón, M F González-Escribano, A Balsa, B Fernández-Gutierrez, I González-Alvaro, M A González-Gay, J Martín