



OPEN ACCESS

EXTENDED REPORT

Genome-wide association study of clinically defined gout identifies multiple risk loci and its association with clinical subtypes

Hirotaka Matsuo,¹ Ken Yamamoto,² Hirofumi Nakaoka,³ Akiyoshi Nakayama,^{1,4} Masayuki Sakiyama,^{1,5} Toshinori Chiba,¹ Atsushi Takahashi,⁶ Takahiro Nakamura,^{6,7} Hiroshi Nakashima,⁸ Yuzo Takada,⁹ Inaho Danjoh,^{10,11} Seiko Shimizu,¹ Junko Abe,¹ Yusuke Kawamura,¹ Sho Terashige,¹ Hiraku Ogata,¹ Seishiro Tatsukawa,¹ Guang Yin,^{12,13} Rieko Okada,¹² Emi Morita,¹² Mariko Naito,¹² Atsumi Tokumasu,¹⁴ Hiroyuki Onoue,¹⁵ Keiichi Iwaya,¹⁶ Toshimitsu Ito,¹⁷ Tappei Takada,¹⁸ Katsuhisa Inoue,¹⁹ Yukio Kato,²⁰ Yukio Nakamura,¹⁰ Yutaka Sakurai,⁸ Hiroshi Suzuki,¹⁸ Yoshikatsu Kanai,²¹ Tatsuo Hosoya,^{22,23} Nobuyuki Hamajima,²⁴ Ituro Inoue,³ Michiaki Kubo,²⁵ Kimiyoshi Ichida,^{22,26} Hiroshi Ooyama,¹⁴ Toru Shimizu,²⁷ Nariyoshi Shinomiya¹

Handling editor Tore K Kvien

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2014-206191>).

For numbered affiliations see end of article.

Correspondence to

Dr Hirotaka Matsuo, Department of Integrative Physiology and Bio-Nano Medicine, National Defense Medical College, 3-2 Namiki, Tokorozawa, Saitama 359-8513, Japan; hmatsuo@ndmc.ac.jp

HM, KY, HNakaoka, AN and MS contributed equally.

Received 1 July 2014
Revised 22 December 2014
Accepted 6 January 2015
Published Online First
2 February 2015



Open Access
Scan to access more
free content



CrossMark

To cite: Matsuo H, Yamamoto K, Nakaoka H, et al. *Ann Rheum Dis* 2016;**75**:652–659.

ABSTRACT

Objective Gout, caused by hyperuricaemia, is a multifactorial disease. Although genome-wide association studies (GWASs) of gout have been reported, they included self-reported gout cases in which clinical information was insufficient. Therefore, the relationship between genetic variation and clinical subtypes of gout remains unclear. Here, we first performed a GWAS of clinically defined gout cases only.

Methods A GWAS was conducted with 945 patients with clinically defined gout and 1213 controls in a Japanese male population, followed by replication study of 1048 clinically defined cases and 1334 controls.

Results Five gout susceptibility loci were identified at the genome-wide significance level ($p < 5.0 \times 10^{-8}$), which contained well-known urate transporter genes (*ABCG2* and *SLC2A9*) and additional genes: rs1260326 ($p = 1.9 \times 10^{-12}$; OR=1.36) of *GCKR* (a gene for glucose and lipid metabolism), rs2188380 ($p = 1.6 \times 10^{-23}$; OR=1.75) of *MYL2-CUX2* (genes associated with cholesterol and diabetes mellitus) and rs4073582 ($p = 6.4 \times 10^{-9}$; OR=1.66) of *CNIH-2* (a gene for regulation of glutamate signalling). The latter two are identified as novel gout loci. Furthermore, among the identified single-nucleotide polymorphisms (SNPs), we demonstrated that the SNPs of *ABCG2* and *SLC2A9* were differentially associated with types of gout and clinical parameters underlying specific subtypes (renal underexcretion type and renal overload type). The effect of the risk allele of each SNP on clinical parameters showed significant linear relationships with the ratio of the case–control ORs for two distinct types of gout ($r = 0.96$ [$p = 4.8 \times 10^{-4}$] for urate clearance and $r = 0.96$ [$p = 5.0 \times 10^{-4}$] for urinary urate excretion).

Conclusions Our findings provide clues to better understand the pathogenesis of gout and will be useful for development of companion diagnostics.

INTRODUCTION

Gout is a common disease caused by deposition of monosodium urate (MSU) crystal due to hyperuricaemia.¹ Humans have long suffered from gout as reported by Hippocrates 2500 years ago.² There have been many famous patients with gout such as Sir Isaac Newton³ in the more recent past, and the numbers are still growing. From the pathophysiological point of view, gout can be classified into the renal underexcretion (RUE) type, the renal overload (ROL) type and the combined type based on clinical parameters⁴ (see online supplementary figure S1).

So far the genome-wide association studies (GWASs) of serum uric acid (SUA) level^{5–16} have identified a number of genetic loci including *SLC2A9* (also known as *GLUT9*) and *ABCG2* (also known as *BCRP*), and subsequent genetic and functional studies have revealed the biological and pathophysiological significance of *ABCG2*.^{4 17 18} Previous GWASs of gout reported a significant association with single-nucleotide polymorphisms (SNPs) of *ABCG2*, *SLC2A9* with European ancestries,^{14 15} and of *ALDH16A1* with Icelanders,¹⁴ while another study with African-American and European ancestries reported no significantly associated SNPs of gout.¹³ All of these studies were, however, performed with cases including self-reported patients with gout, in which clinical information was insufficient. Therefore, the relation to genetic heterogeneity underlying gout subtypes is also unclear. To better understand its genetic basis, we first performed a GWAS of clinically defined gout cases only. We then investigated the relationship between genetic variation and clinical types of gout.

METHODS

Subjects

In the present study, we avoided use of self-reported gout cases and collected only clinically defined gout

cases. All gout cases were clinically diagnosed as primary gout according to the criteria established by the American College of Rheumatology.¹⁹ All patients were assigned from among the Japanese male outpatients at the gout clinics of Midorigaoka Hospital (Osaka, Japan), Kyoto Industrial Health Association (Kyoto, Japan) or Ryougoku East Gate Clinic (Tokyo, Japan). Patients with inherited metabolism disorders including Lesch-Nyhan syndrome were excluded. Finally, 1994 male gout cases were registered as valid case participants. As controls, 2547 individuals were assigned from among Japanese men with normal SUA level (≤ 7.0 mg/dL) and no gout history, who were obtained from BioBank Japan^{11 20} and Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study).²¹

Genotyping and quality control

Genome-wide genotyping was performed with Illumina HumanOmniExpress v1.0 (Illumina) in 946 cases and 1213 controls. Detailed methods of genotyping and quality control are shown in the online supplementary methods and figure S2. Finally, 570 442 SNPs passed filters for 945 cases and 1213 controls.

In total, 123 SNPs passing the significance threshold at $p < 1.0 \times 10^{-5}$ in the GWAS stage were used for subsequent analyses. Among these SNPs, we examined their linkage disequilibrium (LD) and selected 16 SNPs for replication study (see online supplementary methods). These 16 SNPs were then genotyped by an allelic discrimination assay (Custom TaqMan Assay and By-Design, Applied Biosystems) with a LightCycler 480 (Roche Diagnostics).¹⁸ After quality control, subsequent statistical analysis was performed with 1048 cases and 1334 controls.

Statistical analyses for GWAS

We conducted an association analysis using a 2×2 contingency table based on the allele frequency, and p value of association was assessed by χ^2 test. The quantile–quantile plot and the genomic inflation factor were used to assess the presence of systematic bias in the test statistics due to potential population stratification (see online supplementary methods and figure S3).

We then combined results from the GWAS and replication stages by meta-analysis.²² The inverse-variance fixed-effects model meta-analysis was used for estimating summary OR. Cochran's Q test²³ and I^2 statistic^{24 25} were examined to assess heterogeneity in ORs between GWAS and replication study. If heterogeneity was present by the statistical test ($p_{\text{het}} < 0.05$) or measurement ($I^2 > 50\%$), we implemented DerSimonian and Laird random-effects model meta-analysis.²⁶ All the meta-analyses were performed using the STATA V.11.0. Genome-wide significance threshold was set to be $\alpha = 5.0 \times 10^{-8}$ to claim evidence of a significant association. Detailed methods of imputation and per cent variance are shown in the online supplementary methods.

Subtype analyses

Gout contains two distinct types, 'ROL' type and 'RUE' type. The ROL type was defined when urinary urate excretion (UUE) was over $25.0 \text{ mg/h}/1.73 \text{ m}^2$ ($600 \text{ mg/day}/1.73 \text{ m}^2$)^{4 27–29} and their urate clearance (urate clearance/creatinine clearance ratio, FE_{UA}) was 5.5% or over. Also, the RUE type was determined when UUE was $25.0 \text{ mg/h}/1.73 \text{ m}^2$ or under and FE_{UA} was under 5.5%.^{4 30 31} Detailed methods of subtype analyses are described in the online supplementary methods.

RESULTS

Genome-wide association study

Clinical characteristics of participants in this study are shown in online supplementary tables S1–S3. GWAS with 945 clinically defined gout cases and 1213 controls identified SNPs in three loci showing evidence of associations at the genome-wide significance level ($p < 5.0 \times 10^{-8}$): rs2728125 of *ABCG2* ($p = 1.5 \times 10^{-27}$; OR=2.05), rs3775948 of *SLC2A9* ($p = 6.7 \times 10^{-15}$; OR=1.64) and rs2188380 of *MYL2-CUX2* ($p = 5.7 \times 10^{-13}$; OR=1.78, figures 1 and 2, table 1 and online supplementary figure S4).

Replication study was conducted with 1048 cases and 1334 controls. As a result, the three SNPs surpassing the genome-wide significance threshold in the GWAS stage were successfully replicated; rs2728125 ($p = 8.3 \times 10^{-29}$; OR=2.03), rs3775948 ($p = 7.6 \times 10^{-14}$; OR=1.57) and rs2188380 ($p = 2.0 \times 10^{-12}$;




Figure 1 Manhattan plot of a genome-wide association analysis of gout. X-axis shows chromosomal positions. Y-axis shows $-\log_{10} p$ values. The upper and lower dotted lines indicate the genome-wide significance threshold ($p = 5.0 \times 10^{-8}$) and the cut-off level for selecting single-nucleotide polymorphisms for replication study ($p = 1.0 \times 10^{-5}$), respectively.




Figure 2 Regional association plots for six discovered loci of gout. Five regions exceeding the genome-wide significance level (A–E) and one region showing a suggestive association (F). The highest association signal in each panel is located on *ABCG2* (A), *SLC2A9* (B), *MYL2-CUX2* (C), *GCKR* (D), *CNIH-2* (E) and *MAP3K11* (F). Region within 250 kb from single-nucleotide polymorphism (SNP) showing lowest p value is displayed. (Top panel) Plots of $-\log_{10}$ p values for the test of SNP association with gout in the genome-wide association study stage. SNP showing the lowest p value is depicted as a pink diamond. Other SNPs are colour-coded according to the extent of linkage disequilibrium (measured in r^2) with SNP showing the lowest p value. (Middle panel) Recombination rates (centimorgans per Mb) estimated from HapMap Phase II data are plotted. (Bottom panel) RefSeq genes. Genomic coordinates are based on Genomic Reference Consortium GRCh37.

OR=1.73). Additionally, two SNPs showed significant associations at $p < 3.1 \times 10^{-3}$ ($=0.05/16$) with Bonferroni correction; rs1260326 of *GCKR* ($p = 2.8 \times 10^{-6}$; OR=1.32) and rs4073582 of *CNIH-2* ($p = 1.6 \times 10^{-4}$; OR=1.55) as shown in table 1 and online supplementary table S4.

All five SNPs that showed significant associations in the replication study achieved genome-wide significance in the meta-analysis of GWAS and replication study (table 1): rs2728125 ($p_{\text{meta}} = 7.2 \times 10^{-54}$; OR=2.04), rs3775948 ($p_{\text{meta}} = 5.5 \times 10^{-27}$; OR=1.61), rs2188380 ($p_{\text{meta}} = 1.6 \times 10^{-23}$; OR=1.75), rs1260326 ($p_{\text{meta}} = 1.9 \times 10^{-12}$; OR=1.36) and rs4073582 ($p_{\text{meta}} = 6.4 \times 10^{-9}$; OR=1.66). In addition, an intronic SNP of *MAP3K11* (rs10791821) showed a suggestive level of association ($p_{\text{meta}} = 1.0 \times 10^{-7}$; OR=1.57). There was $>80\%$ power to detect a risk variant for OR=1.6 at the genome-wide significance level ($p < 5.0 \times 10^{-8}$) for an SNP with a minor allele frequency of 0.35 (see online supplementary table S5). Imputation was also performed with the GWAS genotyping data for 1 Mb across the identified SNPs of novel loci (rs2188380 of *MYL2-CUX2*, rs1260326 of *GCKR*, rs4073582 of *CNIH-2* and rs10791821 of *MAP3K11*). SNPs that passed the significant threshold of GWAS stage ($p < 1.0 \times 10^{-5}$) in this imputation are shown in online supplementary table S6A–D.

Two dysfunctional SNPs of *ABCG2*

We previously demonstrated that two dysfunctional SNPs of *ABCG2*, rs72552713 (Gln126Ter) and rs2231142 (Gln141Lys), were located on different haplotypes^{4 18} and strongly associated with hyperuricaemia and gout.^{4 18 32} Therefore, we additionally performed genotyping of these two SNPs by an

allelic discrimination assay because SNPs are not on Illumina HumanOmniExpress V1.0 (Illumina). SNP showing the highest significance in the present GWAS (rs2728125) was in strong LD with rs2231142 ($r^2 = 0.76$) but not in LD with rs72552713 ($r^2 = 0.03$). A multivariate logistic regression analysis including these three SNPs of *ABCG2* showed that rs2728125 no longer had a significant association ($p = 0.19$), but rs72552713 and rs2231142, that is, two non-synonymous SNPs, remained highly significant (see online supplementary table S7A, B), indicating that rs2728125 was merely a surrogate marker for rs2231142. Therefore, we used these two non-synonymous variants for subsequent analyses.

Cumulative effect of risk alleles for gout

Accumulation of the number of risk alleles of the gout-associated SNPs (rs3775948, rs2188380, rs1260326, rs4073582, rs72552713 and rs2231142) increased the probability of gout logarithmically. When setting the reference category as having four or fewer risk alleles, ORs for having 5, 6, 7, 8 and 9 or more risk alleles were 1.79 ($p = 3.5 \times 10^{-3}$), 3.16 ($p = 2.3 \times 10^{-10}$), 5.10 ($p = 9.7 \times 10^{-21}$), 10.1 ($p = 5.3 \times 10^{-39}$) and 18.6 ($p = 3.6 \times 10^{-45}$), respectively (see online supplementary figure S5 and table S8).

Subtype analysis of gout

We examined type-specific ORs and the case–subtype heterogeneity test.³³ The subgroup analysis (table 2) showed that the associations of two non-synonymous SNPs of *ABCG2* (rs72552713 and rs2231142) were stronger for the ROL type (ORs=4.35 and 3.37, respectively) than for the RUE type (ORs=1.28 and

Table 1 Five SNPs showing significant association at genome-wide significance level and one suggestive SNP

SNP [¶]	Chromosome	Position (bp) ^{††}	Gene	A1/A2 ^{‡‡}	GWAS†			Replication study ^{‡‡}			Meta-analysis ^{§§}			
					Cases	Controls	OR (95% CI)	p Value	Cases	Controls	OR (95% CI)	p Value	OR (95% CI)	p Value
					Freq.			Freq.			Freq.			
rs2728125	4	89 001 893	ABCG2	C/T	0.40	0.25	2.05 (1.80 to 2.34)	1.5×10 ⁻²⁷	0.40	0.24	2.03 (1.79 to 2.30)	8.3×10 ⁻²⁹	2.04 (1.86 to 2.23)	7.2×10 ⁻⁵⁴
rs3775948	4	9 995 182	SLC2A9	G/C	0.68	0.56	1.64 (1.45 to 1.86)	6.7×10 ⁻¹⁵	0.67	0.56	1.57 (1.40 to 1.77)	7.6×10 ⁻¹⁴	1.61 (1.47 to 1.75)	5.5×10 ⁻²⁷
rs2188380	12	111 386 127	MYL2-CUX2	T/C	0.85	0.76	1.78 (1.52 to 2.08)	5.7×10 ⁻¹³	0.86	0.78	1.73 (1.48 to 2.02)	2.0×10 ⁻¹²	1.75 (1.57 to 1.96)	1.6×10 ⁻²³
rs1260326	2	27 730 940	GCKR	T/C	0.62	0.54	1.39 (1.23 to 1.57)	1.2×10 ⁻⁷	0.61	0.55	1.32 (1.18 to 1.49)	2.8×10 ⁻⁶	1.36 (1.25 to 1.48)	1.9×10 ⁻¹²
rs4073582	11	66 050 712	CNIH-2	G/A	0.95	0.91	1.78 (1.39 to 2.29)	5.3×10 ⁻⁶	0.94	0.91	1.55 (1.23 to 1.96)	1.6×10 ⁻⁴	1.66 (1.40 to 1.96)	6.4×10 ⁻⁹
rs10791821 ^{**}	11	65 368 323	MAP3K11	G/A	0.94	0.90	1.75 (1.38 to 2.22)	2.8×10 ⁻⁶	0.94	0.92	1.41 (1.12 to 1.77)	3.4×10 ⁻³	1.57 (1.33 to 1.85)	1.0×10 ⁻⁷

[†]945 gout cases and 1213 controls.

[‡]1048 gout cases and 1334 controls.

[§]Meta-analyses of the combined GWAS and replication samples (1993 gout cases and 2547 controls).

[¶]dbSNP ID number. A suggestive SNP is marked with †††.

^{‡‡}SNP positions are based on the National Center for Biotechnology Information human genome reference sequence build 37.4.

^{‡‡‡}A1 is a risk-associated allele and A2 is a non-risk-associated allele.

^{§§}Frequency of A1: GWAS, genome-wide association study; SNP, single-nucleotide polymorphism.

1.88, respectively). The differences in ORs between the gout types were highly significant ($p=2.4\times 10^{-5}$ and 1.0×10^{-7} , respectively). The association of rs3775948 of *SLC2A9* was stronger for the RUE type (OR=1.94) than for the ROL type (OR=1.38). The case-subtype heterogeneity test showed a significant difference in ORs ($p=2.7\times 10^{-4}$). The other SNPs evidenced no significant differences. Then, associations between SNPs and clinical parameters (FE_{UA} and UUE) were assessed. Only SNPs that showed a significant difference in ORs between different gout types were significantly associated with FE_{UA} and UUE (table 2 and online supplementary figure S6, table S9); the gout risk alleles of *ABCG2* and *SLC2A9* were associated with increased and decreased levels of these parameters, respectively. The effect of the risk allele of each SNP on clinical parameters showed significant linear relationships with OR in the case-subtype heterogeneity test, which was an estimate of the ratio of the case-control ORs for the gout types ($r=0.96$ [$p=4.8\times 10^{-4}$] for FE_{UA} and $r=0.96$ [$p=5.0\times 10^{-4}$] for UUE) (figure 3).

DISCUSSION

Through the GWAS with clinically defined cases, we identified five gout-associated loci that showed different association patterns in subtype analysis. Previous GWASs of SUA⁵⁻¹⁶ showed genome-wide significant associations with *ABCG2*, *SLC2A9* and *GCKR*. These genes were also reported to have significant associations with gout as a consequence of hyperuricaemia.¹³⁻¹⁵ The present study revealed for the first time that three loci (*GCKR*, *MYL2-CUX2* and *CNIH-2*) were associated with gout at the genome-wide significance level. In particular, *MYL2-CUX2* and *CNIH-2* are novel loci for gout.

The total variance explained by the seven SNPs was estimated to be 9.0% (see online supplementary methods): three SNPs of well-known urate transporter genes (*SLC2A9* and *ABCG2*) with large effects accounted for 6.9%, and the four SNPs identified in this GWAS with modest effects explained 2.1%. Additional discoveries of unidentified genetic variants by performing a meta-analysis of GWAS data sets will improve the explained genetic variation of gout.

ABCG2 and *SLC2A9* are well-known urate transporter genes for urate excretion^{17 18} and renal urate reabsorption,^{34 35} respectively. *ABCG2* is identified to have an association with SUA levels by recent GWASs.⁹⁻¹⁶ Subsequent genetic and functional analysis^{17 18} revealed that *ABCG2* is a high-capacity urate exporter and shows the reduced transport of urate by a common half-functional variant, rs2231142 (Gln141Lys). We also demonstrated that common dysfunctional genotype combinations of *ABCG2* gene (non-functional rs72552713 [Gln126Ter] and rs2231142) are a major cause of hyperuricaemia and gout,¹⁸ especially for early-onset gout.³² We earlier found that the risk alleles of these two SNPs reside on different haplotypes,^{4 18} indicating independent risks of gout. Recently, these dysfunctional SNPs were revealed to decrease extrarenal (intestinal) urate excretion and to cause ROL hyperuricaemia,⁴ through studies with hyperuricaemic patients⁴ and *Abcg2*-knockout mice.^{4 36} This is consistent with the fact that *ABCG2* exporter is expressed on the apical membrane in several tissues, including intestine³⁷ and kidney,³⁸ which have urate-excreting functions in humans.

SLC2A9 is a member of the glucose transporter (GLUT) family. *SLC2A9* was found to transport urate,^{7 34} and several GWAS have demonstrated an association of *SLC2A9* with SUA levels.⁵⁻¹⁶ *SLC2A9* has two isoforms, GLUT9L (long isoform) and GLUT9S (short isoform),³⁴ and is highly expressed in the kidney proximal tubules in humans.³⁹ Genetic and functional

Table 2 Associations of seven SNPs with gout types

SNP†	Gene	Freq.		ROL type vs controls*		RUE type vs controls*		Case-subtype heterogeneity test	
		ROL type	RUE type	OR (95% CI)	p Value	OR (95% CI)	p Value	OR (95% CI)	p Value‡
rs3775948	<i>SLC2A9</i>	0.62	0.70	1.38 (1.14 to 1.68)	1.0×10 ⁻³	1.94 (1.63 to 2.31)	1.0×10 ⁻¹³	0.66 (0.53 to 0.83)	2.7×10 ⁻⁴
rs2188380	<i>MYL2-CUX2</i>	0.84	0.85	1.45 (1.11 to 1.89)	6.5×10 ⁻³	1.47 (1.16 to 1.86)	1.2×10 ⁻³	0.92 (0.68 to 1.25)	0.60
rs1260326§	<i>GCKR</i>	0.60	0.62	1.25 (1.04 to 1.50)	0.016	1.35 (1.15 to 1.58)	3.0×10 ⁻⁴	0.94 (0.77 to 1.14)	0.51
rs4073582	<i>CNIH-2</i>	0.95	0.94	1.96 (1.30 to 2.95)	1.2×10 ⁻³	1.51 (1.09 to 2.08)	0.013	1.26 (0.80 to 1.99)	0.32
rs10791821	<i>MAP3K11</i>	0.93	0.95	1.37 (0.96 to 1.96)	0.084	1.79 (1.26 to 2.54)	1.2×10 ⁻³	0.79 (0.51 to 1.23)	0.30
rs72552713§	<i>ABCG2</i>	0.067	0.029	4.35 (2.82 to 6.72)	3.0×10 ⁻¹¹	1.28 (0.78 to 2.12)	0.32	2.90 (1.77 to 4.75)	2.4×10 ⁻⁵
rs2231142§	<i>ABCG2</i>	0.50	0.38	3.37 (2.76 to 4.12)	2.8×10 ⁻³²	1.88 (1.58 to 2.24)	2.5×10 ⁻¹²	1.76 (1.43 to 2.17)	1.0×10 ⁻⁷

*We performed multivariate logistic regression analyses, in which all seven SNPs, alcohol drinking and body mass index were included in the model. In total, 1613 patients with gout and 1334 controls with genotypes for rs72552713 and rs2231142 of *ABCG2*, which were not on the Illumina OmniExpress platform, were used. Also, 375 and 509 patients with gout were grouped into ROL type and RUE type, respectively.

†dbSNP rs number.

‡p Values <0.05 are shown in bold.

§Non-synonymous SNPs (rs1260326, Leu446Pro; rs72552713, Gln126Ter; and rs2231142, Gln141Lys).

¶Freq., frequency of risk-associated allele; ROL, renal overload; RUE, renal underexcretion; SNP, single-nucleotide polymorphism.

analysis^{34 35} with patients with renal hypouricaemia (RHUC) revealed that RHUC is caused by dysfunctional mutations in *SLC2A9*, which decrease urate reabsorption in the renal proximal tubules. For example, non-functional mutations of either GLUT9L (Arg198Cys and Arg380Trp) or GLUT9S (Arg169Cys and Arg351Trp, corresponding to Arg198Cys and Arg380Trp in GLUT9L), which were found from patients with RHUC, dramatically reduced the urate transport activity.³⁴ Therefore, *SLC2A9* plays an important role in renal urate reabsorption.³⁴ Thus, *SLC2A9* is a causative gene for RHUC type 2,^{34 40} which was confirmed by the report of homozygous mutations in patients with RHUC type 2.³⁵ In our subtype analysis, OR of RUE type was higher than that of ROL type (OR=1.94 and 1.38, respectively, **table 2**), which is compatible with the fact that *SLC2A9* is a transporter for urate reabsorption in human kidney.

Glucokinase regulatory protein (GCKR) controls the activity of glucokinase, which is a major glucose sensor for insulin secretion. GCKR regulates the first step of glycolysis, the phosphorylation of glucose to glucose-6-phosphate.^{41 42} Glucokinase activity is controlled by GCKR, which binds to glucokinase and suppresses its function in the postabsorptive phase. On the other hand, this binding is loosened in the postprandial phase, so that glucokinase could adopt the glycolysis.⁴³ So far, the gout risk allele of rs1260326 (Leu446Pro) has been reported to be associated with lower fasting glucose levels, and inversely, higher levels of triglyceride⁴³⁻⁴⁵ and SUA.^{10 12 15} An association of *GCKR* with dyslipidaemia has also been reported.⁴⁶

MYL2 encodes a regulatory light chain associated with cardiac myosin β (or slow) heavy chain. *MYL2* mutations are associated with mid-left ventricular-type hypertrophic cardiomyopathy. In addition, its association with high-density lipoprotein




Figure 3 Relationships between effects of risk alleles on clinical parameters and ORs in case-subtype heterogeneity test. (A) FE_{UA} and (B) urinary urate excretion (UUE). The seven single-nucleotide polymorphisms (SNPs) listed in **table 2** were examined. OR in case-subtype heterogeneity test is an estimate of the ratio of the case-control OR for the renal overload (ROL) type to that for the renal underexcretion (RUE) type. If an SNP has a stronger effect for the ROL type than for the RUE type, it takes a value >1. Diamonds and lines represent point estimates and their 95% CIs. Pearson's correlation coefficient (r) between the effect on clinical parameters and natural logarithm of OR in case-subtype heterogeneity test and its significance were examined. FE_{UA}, fractional excretion of urate clearance.

cholesterol metabolism was previously reported.⁴⁷ *CUX2* regulates cell-cycle progression⁴⁸ and plays important roles in neural progenitor development in the central nervous system.^{48–49} Its association with type 1 diabetes has also been reported.⁵⁰ Thus, rs2188380 of *MYL2-CUX2* showed an association with gout because *MYL2* and *CUX2* might influence such metabolic pathways. Rs2188380 locates near rs653178 of *ATXN2* (see online supplementary figure S7), which was reported by Köttgen *et al*¹⁵ to have an association with SUA. Rs653178 is, however, monomorphic in the Japanese population of the HapMap project,⁵¹ and we also confirmed it in our samples by genotyping >250 replication cases. Conversely, rs2188380 of *MYL2-CUX2* is monomorphic in European and African populations,⁵¹ while rs2188380 is a common variant in the Japanese population (table 1). Therefore, this SNP was identified as a novel locus of gout in the present study. The differences in study populations could be one of the reasons why rs2188380 was not found in a large European-driven GWAS on urate and gout.¹⁵ Further analyses including fine mapping and functional analysis are required in this region.

CNIH-2 regulates the function of glutamate receptors of the AMPA-subtype assembly at the cell surface of various neurons and glial cells.^{52–53} *CNIH-2* modulates AMPA receptor gating by increasing its cell surface expression. The newly identified rs4073582 of *CNIH-2* was in strong LD with rs801733 in *PACS1* ($r^2=0.97$, figure 2E and see online supplementary figure S4E), which is reported to be associated with severe obesity.⁵⁴ Accordingly, *PACS1* can also be a good candidate for a gout susceptibility gene. Additional genetic dissection and functional analysis will be needed to determine whether these genes or others could play roles with true causality at this locus. Since Okada *et al*¹⁶

previously reported the association between SUA and rs504915 of *NRXN2*, which is near *CNIH-2* and *MAP3K11*, we examined their relationships. They are not in strong LD (see online supplementary table S10), and the association of rs4073582 and rs10791821 remained significant after adjustment with rs504915 (see online supplementary table S11). Therefore, rs4073582 of *CNIH-2*, rs10791821 of *MAP3K11* and rs504915 of *NRXN2* are revealed to be independent of each other.

MAP3K11, also known as mixed lineage kinase 3 (MLK3), is a MAP kinase member and plays a significant role in the activation of c-Jun N-terminal kinase (JNK), a stress-activated protein kinase.⁵⁵ Signalling from the small GTP-binding proteins Rac1 and Cdc42 induces MLK3 to activate the MEKK-SEK-JNK kinase cascade. Interestingly, the JNK pathway is activated when monocytes/macrophages phagocytose MSU crystals,⁵⁶ which cause gouty arthritis. The SNP rs10791821 of *MAP3K11* has been associated with the expression level of *MAP3K11* in monocytes,⁵⁷ and therefore, is likely to be a regulatory SNP. However, further study is required to confirm precise involvement of *MAP3K11* in the development of gout.

Other genes (*CCDC63*, *C2orf16*, *ZNF512*, *RAB1B*, *EHBP1L1* and *KCNK7*) near each of the novel loci, which are found by imputation analysis (see online supplementary table S6A–D), could also be candidate genes of gout, and further studies including functional analyses are warranted.

Most of the gout-related genes are also associated with SUA.¹⁵ In the present study design, to identify novel gout risk loci, clinically defined gout and normouricaemic controls were recruited. Therefore, further investigations with different study designs will be needed to identify gout loci associated with crystal deposition and inflammation.




Figure 4 Differential effects by risk allele on clinical parameters and gout. (A) The risk alleles of *ABCG2* increase UUE and FE_{UA}, which leads to the overloading effect on renal urate excretion and increases the risk of the ROL-type gout. Therefore, patients with risk alleles for the ROL-type gout would be given urate synthesis inhibitors. (B) The risk allele of *SLC2A9* reduces UUE and FE_{UA}, which reflects a decreased renal urate excretion, thereby increasing the risk of the RUE-type gout. Patients with risk alleles for the RUE-type gout would be administered uricosuric agents. FE_{UA}, fractional excretion of urate clearance; ROL, renal overload; RUE, renal underexcretion; SUA, serum uric acid; UUE, urinary urate excretion.

We further investigated the cumulative effect of risk alleles of the five significant loci (*ABCG2*, *SLC2A9*, *MYL2-CUX2*, *GCKR* and *CNIH-2*) on gout risk. The result showed that individuals with five or more risk alleles had a higher risk for gout compared with those having four or fewer risk alleles. The more risk alleles in an individual, the higher became the risk of gout.

Furthermore, the relationship between genetic variation and clinical types of gout was investigated. The results of subtype analyses (table 2, figure 3 and online supplementary figure S6, table S9) indicate that the alleles closely associated with the risk of specific gout type represented differential effects on clinical parameters (FE_{UA} and UUE). This allows the estimation of disturbed urate excretion pathways. An increase of FE_{UA} and UUE by the risk alleles of *ABCG2* leads to the overloading effect on renal urate excretion and causes the ROL-type gout (figure 4A). These estimations are consistent with our previous finding obtained from *Abcg2*-knockout mouse models and hyperuricaemic patients.⁴ In contrast, the reduction of FE_{UA} and UUE by the risk allele of *SLC2A9* reflects a decreased renal urate excretion, thereby increasing the risk of the RUE-type gout (figure 4B). The present study demonstrated that the combination of GWAS of patients with clinically defined gout with actual clinical data is an effective method to analyse genetic heterogeneity among different types of gout.

In summary, we conducted the first GWAS using patients with clinically defined gout only and identified five loci containing two novel loci. Moreover, identified SNPs showed differential effects on different gout types and affected clinical parameters underlying specific types. Thus, genetic testing for gout may well be introduced into future companion diagnostics. For example, patients with risk alleles for ROL-type gout would be given urate synthesis inhibitors^{31 58} such as allopurinol and febuxostat, while patients with risk alleles for RUE-type gout would be administered uricosuric agents^{31 58} including benzbromarone and lesinurad, a selective uric acid reabsorption inhibitor that has just finished its phase III study.^{59 60} Exploring genetic heterogeneity among different gout types will deepen understanding of the aetiology of gout and serve to categorise patients for future personalised treatment.

Author affiliations

- ¹Department of Integrative Physiology and Bio-Nano Medicine, National Defense Medical College, Tokorozawa, Saitama, Japan
- ²Department of Medical Chemistry, Kurume University School of Medicine, Kurume, Fukuoka, Japan
- ³Division of Human Genetics, Department of Integrated Genetics, National Institute of Genetics, Mishima, Shizuoka, Japan
- ⁴Medical Group, Headquarters, Iwo-to Air Base Group, Japan Air Self-Defense Force, Tokyo, Japan
- ⁵Department of Dermatology, National Defense Medical College, Tokorozawa, Saitama, Japan
- ⁶Laboratory for Statistical Analysis, Center for Integrative Medical Sciences, RIKEN, Yokohama, Kanagawa, Japan
- ⁷Laboratory for Mathematics, National Defense Medical College, Tokorozawa, Saitama, Japan
- ⁸Department of Preventive Medicine and Public Health, National Defense Medical College, Tokorozawa, Saitama, Japan
- ⁹The Central Research Institute, National Defense Medical College, Tokorozawa, Saitama, Japan
- ¹⁰Cell Engineering Division, RIKEN BioResource Center, Tsukuba, Ibaraki, Japan
- ¹¹Department of Integrative Genomics, Tohoku Medical Megabank Organization, Tohoku University, Sendai, Miyagi, Japan
- ¹²Department of Preventive Medicine, Nagoya University Graduate School of Medicine, Nagoya, Aichi, Japan
- ¹³Department of Nutritional Sciences, Faculty of Health and Welfare, Seinan Jo Gakuin University, Fukuoka, Japan
- ¹⁴Ryugoku East Gate Clinic, Tokyo, Japan
- ¹⁵Department of Internal Medicine, National Defense Medical College, Tokorozawa, Saitama, Japan

- ¹⁶Department of Pathology, National Defense Medical College, Tokorozawa, Saitama, Japan
- ¹⁷Department of Internal Medicine, Self-Defense Forces Central Hospital, Tokyo, Japan
- ¹⁸Department of Pharmacy, The University of Tokyo Hospital, Tokyo, Japan
- ¹⁹Department of Biopharmaceutics, School of Pharmacy, Tokyo University of Pharmacy and Life Sciences, Tokyo, Japan
- ²⁰Faculty of Pharmacy, Kanazawa University, Kanazawa, Ishikawa, Japan
- ²¹Division of Bio-system Pharmacology, Department of Pharmacology, Graduate School of Medicine, Osaka University, Osaka, Japan
- ²²Division of Kidney and Hypertension, Department of Internal Medicine, Jikei University School of Medicine, Tokyo, Japan
- ²³Department of Pathophysiology and Therapy in Chronic Kidney Disease, Jikei University School of Medicine, Tokyo, Japan
- ²⁴Department of Healthcare Administration, Nagoya University Graduate School of Medicine, Nagoya, Japan
- ²⁵Laboratory for Genotyping Development, Center for Integrative Medical Sciences, RIKEN, Yokohama, Kanagawa, Japan
- ²⁶Department of Pathophysiology, Tokyo University of Pharmacy and Life Sciences, Tokyo, Japan
- ²⁷Midorigaoka Hospital, Osaka, Japan

Acknowledgements We would like to thank all the participants involved in this study. We also thank members of the BioBank Japan Project and Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study) Shizuoka Field for supporting the study. We are indebted to K. Gotanda, Y. Morimoto, N. Katsuta, Y. Utsumi, Y. Kato, H. Sasaki, Y. Takashima, J. Sato, H. Inoue, C. Okada, S. Takeuchi, N. Otani, S. Tomura (National Defense Medical College), T. Tamatsukuri (Jikei University School of Medicine), Y. Oikawa and K. Niwa (Toho University) for genetic analysis; S. Ushida (Ikagaku), H. Fujiwara (Midorigaoka Hospital), A. Hishida and K. Wakai (Nagoya University) for sample collection; M. Hosoyamada, S. Fujimori (Teikyo University), T. Shimizu, T. Sugiura (Kanazawa University), H. Sato (National Defense Medical College), K. Shimono (Toho University) and T. Makino (Nagoya City University) for helpful discussion.

Contributors HM, KY, HNakaoka, AN, MS, TC, II and NS conceived and designed the experiments. ATakahashi, TN, HNaKashima, YT, TT, YS, HS, YKanae, TH and MK assisted with research design. HM, AN, MS, TC, ATokumasu, KIchida, HOoyama and TS collected and analysed clinical data of cases. ATakahashi, GY, RO, EM, MN, NH and MK collected and analysed clinical data of controls. HM, KY, AN, MS, TC, YT, ID, SS, JA, YKawamura, STerashige, HOgata, STatsukawa, YN and NS performed genetic analysis. HNakaoka, ATakahashi, TN, HNaKashima and YS performed statistical analysis. HM, KY, HNakaoka, AN, MS, TC, ATakahashi, TN, HNaKashima, MK, and NS analysed the data. TN, HNaKashima, HOonoue, KIwaya, TI, TT, KIouue, YKato and II provided intellectual input and assisted with the preparation of the manuscript. HM, KY, HNakaoka, AN, MS, TC, ATakahashi and NS wrote the paper.

Funding This work was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan including the MEXT KAKENHI (Grant numbers 221S0002, 25293145, 22689021, 25670307), the Ministry of Health, Labour and Welfare of Japan, the Ministry of Defense of Japan, the Japan Society for the Promotion of Science, the Kawano Masanori Memorial Foundation for Promotion of Pediatrics and the Gout Research Foundation of Japan. The BioBank Japan Project and J-MICC Study (221S0001) were supported by MEXT of Japan.

Competing interests HM, TT and NS have a patent pending based on the work reported in this paper. Other authors have declared that no competing interests exist.

Patient consent Obtained.

Ethics approval All procedures involved in this study were approved by the institutional ethical committees of National Defense Medical College, Nagoya University and RIKEN, and all procedures involved were performed in accordance with the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Kippen I, Klinenberg JR, Weinberger A, *et al.* Factors affecting urate solubility in vitro. *Ann Rheum Dis* 1974;33:313–17.
- 2 Nuki G, Simkin PA. A concise history of gout and hyperuricemia and their treatment. *Arthritis Res Ther* 2006;8(Suppl 1):S1.
- 3 Johnson RJ, Rideout BA. Uric acid and diet—insights into the epidemic of cardiovascular disease. *N Engl J Med* 2004;350:1071–3.

- 4 Ichida K, Matsuo H, Takada T, *et al.* Decreased extra-renal urate excretion is a common cause of hyperuricemia. *Nat Commun* 2012;3:764.
- 5 Li S, Sanna S, Maschio A, *et al.* The GLUT9 gene is associated with serum uric acid levels in Sardinia and Chianti cohorts. *PLoS Genet* 2007;3:e194.
- 6 Döring A, Gieger C, Mehta D, *et al.* SLC2A9 influences uric acid concentrations with pronounced sex-specific effects. *Nat Genet* 2008;40:430–6.
- 7 Vitart V, Rudan I, Hayward C, *et al.* SLC2A9 is a newly identified urate transporter influencing serum urate concentration, urate excretion and gout. *Nat Genet* 2008;40:437–42.
- 8 McArdle PF, Parsa A, Chang YP, *et al.* Association of a common nonsynonymous variant in GLUT9 with serum uric acid levels in old order Amish. *Arthritis Rheum* 2008;58:2874–81.
- 9 Dehghan A, Köttgen A, Yang Q, *et al.* Association of three genetic loci with uric acid concentration and risk of gout: a genome-wide association study. *Lancet* 2008;372:1953–61.
- 10 Kolz M, Johnson T, Sanna S, *et al.* Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. *PLoS Genet* 2009;5:e1000504.
- 11 Kamatani Y, Matsuda K, Okada Y, *et al.* Genome-wide association study of hematological and biochemical traits in a Japanese population. *Nat Genet* 2010;42:210–15.
- 12 Yang Q, Köttgen A, Dehghan A, *et al.* Multiple genetic loci influence serum urate levels and their relationship with gout and cardiovascular disease risk factors. *Circ Cardiovasc Genet* 2010;3:523–30.
- 13 Tin A, Woodward OM, Kao WH, *et al.* Genome-wide association study for serum urate concentrations and gout among African Americans identifies genomic risk loci and a novel URA1 loss-of-function allele. *Hum Mol Genet* 2011;20:4056–68.
- 14 Sulem P, Gudbjartsson DF, Walters GB, *et al.* Identification of low-frequency variants associated with gout and serum uric acid levels. *Nat Genet* 2011;43:1127–30.
- 15 Köttgen A, Albrecht E, Teumer A, *et al.* Genome-wide association analyses identify 18 new loci associated with serum urate concentrations. *Nat Genet* 2013;45:145–54.
- 16 Okada Y, Sim X, Go MJ, *et al.* Meta-analysis identifies multiple loci associated with kidney function-related traits in east Asian populations. *Nat Genet* 2012;44:904–9.
- 17 Woodward OM, Köttgen A, Coresh J, *et al.* Identification of a urate transporter, ABCG2, with a common functional polymorphism causing gout. *Proc Natl Acad Sci U S A* 2009;106:10338–42.
- 18 Matsuo H, Takada T, Ichida K, *et al.* Common defects of ABCG2, a high-capacity urate exporter, cause gout: a function-based genetic analysis in a Japanese population. *Sci Transl Med* 2009;1:5ra11.
- 19 Wallace SL, Robinson H, Masi AT, *et al.* Preliminary criteria for the classification of the acute arthritis of primary gout. *Arthritis Rheum* 1977;20:895–900.
- 20 Nakamura T, Shi D, Tzetzis M, *et al.* Meta-analysis of association between the ASPN D-repeat and osteoarthritis. *Hum Mol Genet* 2007;16:1676–81.
- 21 Hamajima N, J-MICC Study Group. The Japan Multi-Institutional Collaborative Cohort Study (J-MICC Study) to detect gene-environment interactions for cancer. *Asian Pac J Cancer Prev* 2007;8:317–23.
- 22 Nakaoka H, Inoue I. Meta-analysis of genetic association studies: methodologies, between-study heterogeneity and winner's curse. *J Hum Genet* 2009;54:615–23.
- 23 Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954;10:101–29.
- 24 Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- 25 Higgins JP, Thompson SG, Deeks JJ, *et al.* Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- 26 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- 27 Becker MA. Hyperuricemia and gout. In: Scriver CR, Childs B, Kinzler KW, Vogelstein B, eds. *The Metabolic & Molecular Bases of Inherited Disease*. 8th edn. New York: McGraw-Hill, 2001:2513–35.
- 28 Wortmann RL. Disorders of purine and pyrimidine metabolism. In: Fauci AS, Braunwald E, Kasper D, Hauser SL, Long DL, Jameson JL, *et al.*, eds. *Harrison's principles of internal medicine*. 17th edn. New York: McGraw-Hill, 2008:2444–9.
- 29 Wortmann RL. Gout and hyperuricemia. *Curr Opin Rheumatol* 2002;14:281–6.
- 30 Urano W, Taniguchi A, Anzai N, *et al.* Sodium-dependent phosphate cotransporter type 1 sequence polymorphisms in male patients with gout. *Ann Rheum Dis* 2010;69:1232–4.
- 31 The guideline revising committee of the Japanese Society of Gout and Nucleic Acid Metabolism. Diagnosis of hyperuricemia and gout. In: The guideline revising committee of the Japanese Society of Gout and Nucleic Acid Metabolism, ed. *Guideline for the Management of Hyperuricemia and Gout*. 2nd edn. Osaka: Medical Review, 2010:60–72.
- 32 Matsuo H, Ichida K, Takada T, *et al.* Common dysfunctional variants in ABCG2 are a major cause of early-onset gout. *Sci Rep* 2013;3:2014.
- 33 Nakaoka H, Takahashi T, Akiyama K, *et al.* Differential effects of chromosome 9p21 variation on subphenotypes of intracranial aneurysm: site distribution. *Stroke* 2010;41:1593–8.
- 34 Matsuo H, Chiba T, Nagamori S, *et al.* Mutations in glucose transporter 9 gene SLC2A9 cause renal hypouricemia. *Am J Hum Genet* 2008;83:744–51.
- 35 Dinour D, Gray NK, Campbell S, *et al.* Homozygous SLC2A9 mutations cause severe renal hypouricemia. *J Am Soc Nephrol* 2010;21:64–72.
- 36 Hosomi A, Nakanishi T, Fujita T, *et al.* Extra-renal elimination of uric acid via intestinal efflux transporter BCRP/ABCG2. *PLoS ONE* 2012;7:e30456.
- 37 Maliepaard M, Scheffer GL, Faneyte IF, *et al.* Subcellular localization and distribution of the breast cancer resistance protein transporter in normal human tissues. *Cancer Res* 2001;61:3458–64.
- 38 Huls M, Brown CD, Windass AS, *et al.* The breast cancer resistance protein transporter ABCG2 is expressed in the human kidney proximal tubule apical membrane. *Kidney Int* 2008;73:220–5.
- 39 Augustin R, Carayannopoulos MO, Dowd LO, *et al.* Identification and characterization of human glucose transporter-like protein-9 (GLUT9): alternative splicing alters trafficking. *J Biol Chem* 2004;279:16229–36.
- 40 Kawamura Y, Matsuo H, Chiba T, *et al.* Pathogenic GLUT9 mutations causing renal hypouricemia type 2 (RHUC2). *Nucleosides Nucleotides Nucleic Acids* 2011;30:1105–11.
- 41 Brown KS, Kalinowski SS, Megill JR, *et al.* Glucokinase regulatory protein may interact with glucokinase in the hepatocyte nucleus. *Diabetes* 1997;46:179–86.
- 42 Slosberg ED, Desai UJ, Fanelli B, *et al.* Treatment of type 2 diabetes by adenoviral-mediated overexpression of the glucokinase regulatory protein. *Diabetes* 2001;50:1813–20.
- 43 Hishida A, Morita E, Naito M, *et al.* Associations of apolipoprotein A5 (APOA5), glucokinase (GCK) and glucokinase regulatory protein (GCKR) polymorphisms and lifestyle factors with the risk of dyslipidemia and dysglycemia in Japanese—a cross-sectional data from the J-MICC Study. *Endocr J* 2012;59:589–99.
- 44 Vaxillaire M, Cavalcanti-Proenca C, Dechaume A, *et al.* The common P446L polymorphism in GCKR inversely modulates fasting glucose and triglyceride levels and reduces type 2 diabetes risk in the DESIR prospective general French population. *Diabetes* 2008;57:2253–7.
- 45 Shen H, Pollin TI, Damcott CM, *et al.* Glucokinase regulatory protein gene polymorphism affects postprandial lipemic response in a dietary intervention study. *Hum Genet* 2009;126:567–74.
- 46 Dupuis J, Langenber G, Prokopenko I, *et al.* New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* 2010;42:105–16.
- 47 Kim YJ, Go MJ, Hu C, *et al.* Large-scale genome-wide association studies in East Asians identify new genetic loci influencing metabolic traits. *Nat Genet* 2011;43:990–5.
- 48 Lulianella A, Sharma M, Durnin M, *et al.* Cux2 (Cutl2) integrates neural progenitor development with cell-cycle progression during spinal cord neurogenesis. *Development* 2008;135:729–41.
- 49 Franco SJ, Gil-Sanz C, Martinez-Garay I, *et al.* Fate-restricted neural progenitors in the mammalian cerebral cortex. *Science* 2012;337:746–9.
- 50 Huang J, Ellinghaus D, Franke A, *et al.* 1000 Genomes-based imputation identifies novel and refined associations for the Wellcome Trust Case Control Consortium phase 1 data. *Eur J Hum Genet* 2012;20:801–5.
- 51 International HapMap Consortium. The International HapMap Project. *Nature* 2003;426:789–96.
- 52 Schwenk J, Harmel N, Zolles G, *et al.* Functional proteomics identify cornichon proteins as auxiliary subunits of AMPA receptors. *Science* 2009;323:1313–19.
- 53 Herring BE, Shi Y, Suh YH, *et al.* Cornichon proteins determine the subunit composition of synaptic AMPA receptors. *Neuron* 2013;77:1083–96.
- 54 Wheeler E, Huang N, Bochukova EG, *et al.* Genome-wide SNP and CNV analysis identifies common and low-frequency variants associated with severe early-onset obesity. *Nat Genet* 2013;45:513–17.
- 55 Teramoto H, Coso OA, Miyata H, *et al.* Signaling from the small GTP-binding proteins Rac1 and Cdc42 to the c-Jun N-terminal kinase/stress-activated protein kinase pathway. A role for mixed lineage kinase 3/protein-tyrosine kinase 1, a novel member of the mixed lineage kinase family. *J Biol Chem* 1996;271:27225–8.
- 56 Liu R, O'Connell M, Johnson K, *et al.* Extracellular signal-regulated kinase 1/extracellular signal-regulated kinase 2 mitogen-activated protein kinase signaling and activation of activator protein 1 and nuclear factor kappaB transcription factors play central roles in interleukin-8 expression stimulated by monosodium urate monohydrate and calcium pyrophosphate crystals in monocytic cells. *Arthritis Rheum* 2000;43:1145–55.
- 57 Zeller T, Wild P, Szymczak S, *et al.* Genetics and beyond—The transcriptome of human monocytes and disease susceptibility. *PLoS ONE* 2010;5:e10693.
- 58 Mody GM, Tikly M, Kalla AA, *et al.* Approach to arthritis: clinical guideline 2003. *S Afr Med J* 2003;93:949–60.
- 59 AstraZeneca. AstraZeneca announces topline results from phase III monotherapy study of lesinurad in gout patients. [Press release] 13 December 2013. <http://www.astrazeneca.com/Media/Press-releases/Article/13122013-astrazeneca-announces-topline-results-from-phase-iii> (accessed 1 Jul 2014).
- 60 Fleischmann R, Kerr B, Yeh LT, *et al.* Pharmacodynamic, pharmacokinetic and tolerability evaluation of concomitant administration of lesinurad and febuxostat in gout patients with hyperuricaemia. *Rheumatology (Oxford)* 2014;53:2167–74.