

Response to: 'Renal biopsies should be performed whenever treatment strategies depend on renal involvement' by Chemouny *et al*

We thank Chemouny *et al* for their letter and concur with their conclusions.¹ As we state²: "A positive biopsy for ANCA associated vasculitis (AAV) is helpful when considering an initial diagnosis or recurrent disease". In our view, renal biopsy is important to establish diagnosis and may also provide an indication of prognostic trajectory and although existing classification systems need further validation, changes like glomerular sclerosis have obvious adverse prognostic value for patients with AAV.^{3–5} The Delphi process, for the scope of the current recommendations, identified the role of biopsy at both diagnosis and follow-up as an important item for update. Histopathological evidence of vasculitis, such as pauci-immune glomerulonephritis or necrotising vasculitis in any organ, remains the gold standard for diagnostic purposes. The likely diagnostic yield varies and is dependent on the organ targeted and in patients with granulomatosis with polyangiitis (GPA) with renal involvement can be as high as 91.5% from renal biopsy.⁶ As Chemouny *et al* have demonstrated, a renal biopsy was definitive in determining their management decisions. However, during follow-up when relapses occur, it may be prudent to consider judicious use of further kidney biopsy during suspected renal relapse because the cause for acute kidney injury may be due to another cause other than AAV.⁷

M Yates,^{1,2} D R Jayne,³ C Mukhtyar²

¹Norwich Medical School, University of East Anglia, Bob Champion Research and Education Building, Norwich, UK

²Department of Rheumatology, Norfolk and Norwich University Hospital, Norwich, UK

³Lupus and Vasculitis Unit, Addenbrooke's Hospital, Cambridge, UK

Correspondence to Dr M Yates, Norwich Medical School, Bob Champion Research and Education Building, Colney Lane, Norwich NR4 7UY, UK; m.yates@uea.ac.uk

Contributors The authors wrote the response to the eLetter.

Competing interests None declared.

Provenance and peer review Commissioned; internally peer reviewed.



CrossMark

To cite Yates M, Jayne DR, Mukhtyar C. *Ann Rheum Dis* 2017;**76**:e28.

Accepted 3 January 2017

Published Online First 25 January 2017



► <http://dx.doi.org/10.1136/annrheumdis-2016-210933>

Ann Rheum Dis 2017;**76**:e28. doi:10.1136/annrheumdis-2016-210962

REFERENCES

- Chemouny JM, Divard G, Sannier A, *et al*. Renal biopsies should be performed whenever treatment strategies depend on renal involvement. *Ann Rheum Dis* 2017;**76**:e27.
- Yates M, Watts RA, Bajema IM, *et al*. EULAR/ERA-EDTA recommendations for the management of ANCA-associated vasculitis. *Ann Rheum Dis* 2016;**75**:1583–94.
- Berden AE, Ferrario F, Hagen EC, *et al*. Histopathologic classification of ANCA-associated glomerulonephritis. *J Am Soc Nephrol* 2010;**21**: 1628–36.
- Chang DY, Wu LH, Liu G, *et al*. Re-evaluation of the histopathologic classification of ANCA-associated glomerulonephritis: a study of 121 patients in a single center. *Nephrol Dial Transplant* 2012;**27**:2343–9.
- Noone DG, Twilt M, Hayes WN, *et al*. The new histopathologic classification of ANCA-associated GN and its association with renal outcomes in childhood. *Clin J Am Soc Nephrol* 2014;**9**:1684–91.
- Aasarød K, Bostad L, Hammerstrøm J, *et al*. Renal histopathology and clinical course in 94 patients with Wegener's granulomatosis. *Nephrol Dial Transplant* 2001;**16**:953–60.
- Choudhry WM, Nori US, Nadasdy T, *et al*. An unexpected cause of acute kidney injury in a patient with ANCA associated vasculitis. *Clin Nephrol* 2016;**85**:289–95.